arXiv:1503.07949v1 [quant-ph] 27 Mar 2015

Two-channel Optimal Quantum Teleportation

Quan Quan[1], Shao-Ming Fei*[2,3], Heng Fan[4] and Wen-Li Yang*[1,5]

[1]Institute of Modern Physics, Northwest University, Xi'an 710069, China
[2]School of Mathematical Sciences, Capital Normal University, Beijing 100048, China
[3]Max-Planck-Institute for Mathematics in the Sciences, Leipzig 04103, Germany
[4]Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China
[5]Center for Mathematics and Information Interdisciplinary Sciences, Beijing, 100048, China

Abstract

We investigate two-channel scenarios of optimal quantum teleportation based on Bell measurements and GHZ measurements. The detailed protocols are presented and the general expressions of the optimal teleportation fidelities are derived, which turn out to be functions of two-channel fully entangled fractions – invariants under local unitary transformations. We prove that the set of states which are useful for two-channel teleportation is convex and compact. Hence witness operators exist to separate states that are useful for optimal teleportation from the rest ones. Moreover, we show that our two-channel teleportation fidelity is better than the usual one channel ones. The corresponding experimental schemes for two-channel teleportation of photon states are also presented.

Introduction

Quantum teleportation plays an important role in quantum information processing [1], which gives ways to transmit an unknown quantum state from a sender traditionally named "Alice" to a receiver "Bob" who are spatially separated by using classical communication and quantum resources [2, 3, 4, 5]. In [6], the authors considered the one channel optimal teleportation: Alice and Bob previously share a pair of particles in an arbitrary mixed entangled state χ. In order to teleport an unknown state to Bob, Alice first performs a joint Bell measurement on her particles and tells her result to Bob by the classical communication channel. Bob tries his best to choose an particular unitary transformation which depends on the quantum resource χ, so as to get the maximal transmission fidelity. The transmission fidelity of such optimal teleportation is given by the fully entangled fraction (FEF) [7] of the quantum resource. It shows that when the resource χ is a maximally entangled pure state, the corresponding optimal fidelity is equal to 1. However, Alice and Bob usually share a mixed entangled state due to the decoherence, and the fidelity is less than 1. One of the major goals of quantum information theory is to find the optimal ways to make use of noisy channels for communication or establish better entanglement [8].

One way to improve the fidelity of teleportation is to distill entanglement [9], which refers to the procedure of converting mixed entanglement states to singlets by using many copies of the entangled resources. The distillation of pure states is often referred as entanglement concentration [10]. For mixed states, since the distillation protocol presented in [9], fruitful results have been obtained [11, 12, 13, 14]. It has been found that some entangled quantum mixed states, called bound entangled states, are not distillable [15]. The nonlocality and usefulness of bound entangled states in information processing have been extensively studied [16, 17, 18, 19].

The problem to use distillation procedure to improve the teleportation fidelity is that the complicated distillation protocol may have to be repeated for infinitely many times to bring out a singlet. Moreover, in each round the desired results are usually obtained probabilistically, usually with an extremely low possibility to get a expected measurement outcome. Hence a lot of copies of resource states are needed. Above all, if the entangled resources are bound entangled ones, the distillation fails.

In this paper, instead of first doing distillation of two pairs of entangled resources and then doing teleportation by using the distilled pair of entangled resource, we consider to do teleportation directly by using these two pairs of entangled resources. We give two two-channel optimal teleportation protocols, one based on usual two-particle Bell measurement and another one on three-particle GHZ measurement. After tedious algebraic calculations we present the explicit formulae of these teleportation channels. The corresponding optimal teleportation fidelities are derived and analyzed. The two-channel teleportation fidelity is shown to be better than the usual one channel teleportation fidelity. The experimental schemes to demonstrate two-channel teleportation of photon states are also presented.

Two-channel optimal teleportation protocol based on Bell measurements

Let H denote an n-dimensional Hilbert space, with $\{|i\rangle, i = 0, ..., n-1, n < \infty\}$ as orthogonal normalized basis. A set of unitary matrices U_{st} in H can be defined as follows: $U_{st} = h^t g^s$, where h and g are $n \times n$ matrices such that $h|j\rangle = |(j+1)/mod \quad n\rangle$ and $g|j\rangle = w^j|j\rangle$, with $w = exp\{-2\sqrt{-1}\pi/n\}$. One has the following relations [20]: $tr(U_{st}^\dagger U_{s't'}) = n\delta_{tt'}\delta_{ss'}$, $U_{st}U_{st}^\dagger = I_{n\times n}$. The generalized Bell states [6] are given by $|\Phi_{st}\rangle = (1 \otimes U_{st})|\Phi\rangle$, where $|\Phi\rangle = |\Phi_{00}\rangle = \frac{1}{\sqrt{n}}\sum_{i=0}^{n-1}|ii\rangle$ is the maximally entangled pure state. The n^2 generalized Bell states $\{|\Phi_{st}\rangle\} = \frac{1}{\sqrt{n}}\sum_{i,j}(U_{st})_{ij}^*|ij\rangle$ form a complete orthogonal normalized basis of the $H \otimes H$ space. Throughout this paper we adopt the standard notations: for any matrix $A \in \mathrm{End}(H)$, A_α is an embedding operator in the tensor space $H \otimes H \otimes \cdots \otimes H$, which acts as A on the α-th space and as identity on the other spaces; and for any matrix $U \in \mathrm{End}(H \otimes H)$, $U_{\alpha\beta}$ is an embedding operator in $H \otimes H \otimes \cdots \otimes H$, which acts as identity on the spaces except for the α-th and β-th ones.

The two-channel teleportation protocol is as follows. Initially Alice and Bob share two pairs of entangled resources, see Fig. 1. Particles 1 and 2, 3 and 4 are in an entangled state χ, respectively. Particles 1 and 3 are in Alice's side, while particles 2 and 4 are in Bob's side. Alice wants to transmit an unknown state ρ_{in} of particle 0 to Bob. Firstly, Alice (resp. Bob) performs a joint local unitary operation W (resp. V) on particles 1 (resp. 2) and particle 3 (resp. 4). Then she makes a joint Bell measurement on particles 0 and 1, which maps the state of particles 0 and 1 to a basic one, say $|ij\rangle_{01}$, where the subindices denote the spaces with respect to the corresponding particles. She informs Bob the measurement results by classical means. According to these measurement results, Bob chooses a corresponding unitary transformation T_{ij} on particle 2 to achieve the maximal teleportation fidelity. After some tedious calculations, we have:



Figure 1: Scheme of two-channel optimal teleportation protocol based on Bell measurements.

Theorem 1 For any unknown input state ρ_{in}, the teleportation protocol maps the state ρ_{in} to state $\Lambda^{\{T_{st},W,V\}}_{\chi^{\otimes 2}}(\rho_{in})$,

$$\begin{aligned}\Lambda^{\{T_{st},W,V\}}_{\chi^{\otimes 2}}(\rho_{in}) \;=\;& \frac{1}{n^3}\sum_{s_1,t_1,s'_1t'_1}\;\sum_{s_2,t_2,s'_2t'_2}\;\sum_{s,t}\langle\Phi_{s_1t_1}|\chi|\Phi_{s'_1t'_1}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s'_2t'_2}\rangle tr_4[(T_{st})_2V_{24}(U_{s_1t_1})_2\\ &(U_{s_2t_2})_4W_{24}(U_{st})_2^\dagger(\rho_{in})_2(U_{st})_2W_{24}^\dagger(U^\dagger_{s'_1t'_1})_2(U_{s'_2t'_2})_4^\dagger V_{24}^\dagger(T_{st})_2^\dagger].\end{aligned}$$

Proof. First consider that the unknown initial input state ρ_{in} that Alice wants to teleport is a pure state, $|\phi\rangle = \sum_\nu \alpha_\nu|\nu\rangle$.

1. The two entangled resource states are pure: $\chi^{\otimes 2} = |\Psi\rangle\langle\Psi|$, where

$$|\Psi\rangle = \sum_{i,j=0}^{n-1}\sum_{k,l=0}^{n-1} a_{ij}|ij\rangle \otimes a_{kl}|kl\rangle, \qquad \sum_{i,j=0}^{n-1}|a_{ij}|^2 = 1.$$

The initial state is $|\phi\rangle_0|\Psi\rangle_{1234}$. Alice and Bob apply the unitary transformations W and V to their two resource particles respectively. Before the measurement, the initial state becomes $|\phi\rangle_0 W_{13}V_{24}|\Psi\rangle_{1234} = \sum_{i,j=0,k,l=0}^{n-1}\sum_{i',j',k',l',\nu=0}^{n-1} a_{ij}a_{kl}W^{ik}_{i'k'}V^{jl}_{j'l'}\alpha_\nu|\nu i'j'k'l'\rangle_{01234}$.

After Alice's joint Bell measurement based on $|\Phi_{st}\rangle$ on particles 0 and 1, we get: $\langle\Phi_{st}|_{01}(|\phi\rangle_0 W_{13}V_{24}|\Psi\rangle_{1234}) = V_{24}A_2A_4W_{24}(U^\dagger_{st})_2|\phi\rangle_2|\Phi\rangle_{34}$, where A is the $n\times n$ matrix with elements $(A)_{ij} = a_{ij}$. Receiving the Alice's measurement outcomes, correspondingly Bob applies a unitary operator T_{st} on particle 2. The resulting state becomes $(T_{st})_2V_{24}A_2A_4W_{24}(U^\dagger_{st})_2|\phi\rangle_2|\Phi\rangle_{34}$. Taking partial trace over the spaces with respect to particles 1 and 4, we have

$$\begin{aligned}
&\Lambda^{\{T_{st},W,V\}}_{\chi^{\otimes 2}}(\rho_{in})\\
=\;& \sum_{s,t} tr_{34}[(T_{st})_2V_{24}A_2A_4W_{24}(U_{st})^\dagger_2|\phi\rangle_2\langle\phi|_2|\Phi\rangle_{34}\langle\Phi|_{34}(U_{st})_2W^\dagger_{24}A^\dagger_2A^\dagger_4V^\dagger_{24}(T_{st})^\dagger_2]\\
=\;& \sum_{s,t}\frac{1}{n}tr_4[(T_{st})_2V_{24}A_2A_4W_{24}(U_{st})^\dagger_2|\phi\rangle_2\langle\phi|_2(U_{st})_2W^\dagger_{24}A^\dagger_2A^\dagger_4V^\dagger_{24}(T_{st})^\dagger_2]. \qquad (1)
\end{aligned}$$

2. Now consider the case of arbitrary entangled mixed resources,

$$\chi^{\otimes 2} = \sum_{\alpha,\beta} P_\alpha P_\beta|\Psi_{\alpha\beta}\rangle\langle\Psi_{\alpha\beta}|,$$

where

$$|\Psi_{\alpha\beta}\rangle = \sum_{i,j=0}^{n-1}\sum_{k,l=0}^{n-1} a^{(\alpha)}_{ij}|ij\rangle \otimes a^{(\beta)}_{kl}|kl\rangle,$$

$0 \le P_{\alpha(\beta)} \le 1$ and $\sum_{\alpha(\beta)} P_{\alpha(\beta)} = 1$. Similar to the derivation of (1), we have

$$\begin{aligned}
\Lambda^{\{T_{st},W,V\}}_{\chi^{\otimes 2}}(\rho_{in}) \;=\;& \frac{1}{n}\sum_{s,t}\sum_{\alpha,\beta}P_\alpha P_\beta tr_4[(T_{st})_2V_{24}A^{(\alpha)}_2A^{(\beta)}_4W_{24}(U^\dagger_{st})_2|\phi\rangle_2\langle\phi|_2\\
& (U_{st})_2W^\dagger_{24}A^{(\alpha)\dagger}_2A^{(\beta)\dagger}_4V^\dagger_{24}(T_{st})^\dagger_2],
\end{aligned}$$

where $(A)^{(\alpha/\beta)}_{ij} = a^{(\alpha/\beta)}_{ij}$.

Since each matrix $A^{(\alpha)}$ can be decomposed in the basis of U_{st}: $A^{(\alpha)} = \sum_{s,t}a^{(\alpha)}_{st}U_{st}$, by using the relation [6] $n\sum_\alpha p_\alpha a^{(\alpha)}_{st}a^{(\alpha)*}_{s't'} = \langle\Phi_{st}|\chi|\Phi_{s't'}\rangle$, we have

$$\begin{aligned}
\Lambda^{\{T_{st},W,V\}}_{\chi^{\otimes 2}}(\rho_{in}) \;=\;& \frac{1}{n^3}\sum_{s_1,t_1,s'_1t'_1}\;\sum_{s_2,t_2,s'_2t'_2}\;\sum_{s,t}\langle\Phi_{s_1t_1}|\chi|\Phi_{s'_1t'_1}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s'_2t'_2}\rangle tr_4[(T_{st})_2V_{24}(U_{s_1t_1})_2\\
& (U_{s_2t_2})_4W_{24}(U_{st})^\dagger_2|\phi\rangle_2\langle\phi|_2(U_{st})_2W^\dagger_{24}(U^\dagger_{s'_1t'_1})_2(U_{s'_2t'_2})^\dagger_4V^\dagger_{24}(T_{st})^\dagger_2].
\end{aligned}$$

It is straightforward to show that the above relation is valid too for any mixed input state ρ_{in}. $\square$

Remark If we chose $W_{13} = V_{24} = I_{n^2\times n^2}$, $\Lambda_{\chi^{\otimes 2}}^{\{T_{st},W,V\}}(\rho_{in})$ reduces to the one-channel teleportation protocol $\Lambda^{(\chi)}(\{T\})(\rho)$ in [6]:

$$\begin{aligned}\Lambda_{\chi^{\otimes 2}}^{\{T_{st},W,V\}}(\rho_{in}) &= \frac{1}{n^3}\sum_{s_1,t_1,s_1't_1'}\sum_{s_2,t_2,s_2't_2'}\sum_{s,t}\langle\Phi_{s_1t_1}|\chi|\Phi_{s_1't_1'}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s_2't_2'}\rangle \\ &\quad\times \mathrm{tr}_4[(T_{st})_2(U_{s_1t_1})_2(U_{s_2t_2})_4(U_{st})_2^\dagger|\phi\rangle_2\langle\phi|_2(U_{st})_2(U^\dagger_{s_1't_1'})_2(U_{s_2't_2'})_4^\dagger(T_{st})_2^\dagger], \\ &= \Lambda^{(\chi)}(\{T\})(\rho_{in})\times\frac{1}{n}\sum_{s_2,t_2}\sum_{s_2',t_2'}\langle\Phi_{s_2t_2}|\chi|\Phi_{s_2't_2'}\rangle\,\mathrm{tr}[U_{s_2t_2}U^\dagger_{s_2't_2'}] = \Lambda^{(\chi)}(\{T\})(\rho_{in}).\end{aligned}$$

Hence the two-channel protocol is always at least as good as the one-channel one.

Moreover, one can prove that the quantum teleportation channel $\Lambda_{\chi^{\otimes 2}}^{\{T_{st},W,V\}}$ is trace preserving,

$$\begin{aligned}&tr[\Lambda_{\chi^{\otimes 2}}^{\{T_{st},W,V\}}(\rho_{in})] \\ =\ & \frac{1}{n^3}\sum_{s_1,t_1,s_1't_1'}\sum_{s_2,t_2,s_2't_2'}\langle\Phi_{s_1t_1}|\chi|\Phi_{s_1't_1'}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s_2't_2'}\rangle tr_2\{(T_{st})_2(U_{s_1t_1})_2 tr_4[(U_{s_2t_2})_4 \\ & W_{24}W_{24}^\dagger(U_{s_2't_2'})_4^\dagger](U^\dagger_{s_1't_1'})_2(T_{st})_2^\dagger\} \\ =\ & \frac{1}{n^2}\sum_{s_1,t_1,s_1't_1'}\sum_{s_2,t_2,s_2't_2'}\langle\Phi_{s_1t_1}|\chi|\Phi_{s_1't_1'}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s_2't_2'}\rangle tr_2[(U_{s_1t_1})_2(U^\dagger_{s_1't_1'})_2]tr_4[(U_{s_2t_2})_4(U_{s_2't_2'})_4^\dagger] \\ =\ & 1,\end{aligned}$$

where in the first equality we have used the identity $\sum_{s,t}U_{st}^\dagger AU_{st} = ntr(A)I_{n\times n}$, for any $n\times n$ matrix A.

Theorem 2 The optimal teleportaion fidelity of the two-channel protocol $f_{2\max}(\chi)$ is given by

$$f_{2\max}(\chi) = \frac{nF_2(\chi)}{(n+1)} + \frac{1}{n+1}, \tag{2}$$

where $F_2(\chi)$, we call it two-channel fully entangled fraction (TFEF), is given by

$$F_2(\chi) = \max_{\Omega,V\in U(n^2)}\{\langle\Phi|_{12}\,\mathrm{tr}_{34}[\Omega_{13}V_{24}\chi_{12}\chi_{34}\Omega_{13}^\dagger V_{24}^\dagger]|\Phi\rangle_{12}\}. \tag{3}$$

Proof. Let $U(n)$ be an irreducible n-dimensional representation of unitary group G. By using the Schur's lemma

$$\begin{aligned}
&\int_G dg(U^\dagger(g)\otimes U^\dagger(g))\sigma(U(g)\otimes U(g)) = \alpha_1 I\otimes I + \alpha_2 P,\\
&\alpha_1 = \frac{n^2\,\mathrm{tr}(\sigma) - n\,\mathrm{tr}(\sigma P)}{n^2(n^2-1)}, \qquad \alpha_2 = \frac{n^2\,\mathrm{tr}(\sigma P) - n\,\mathrm{tr}(\sigma)}{n^2(n^2-1)},
\end{aligned}$$

where σ is any operator acting on the tensor space, P is the flip operator, dg is the Haar measure on G normalized by $\int_G dg = 1$, we get the fidelity of the two-channel teleportation protocol,

$$\begin{aligned}
f_2(\chi) &= \overline{\langle\phi_{in}|\Lambda^{\{T_{st},W,V\}}_{\chi^{\otimes 2}}(\rho_{in})|\phi_{in}\rangle} = \frac{1}{n^3}\sum_{s_1,t_1}\sum_{s_1',t_1'}\sum_{s_2,t_2}\sum_{s_2',t_2'}\langle\Phi_{s_1t_1}|\chi|\Phi_{s_1't_1'}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s_2't_2'}\rangle\\
&\quad \times\sum_{s,t,j,k}\langle 00|\int_G[U(g)^\dagger\otimes U(g)^\dagger][\langle j|_4(T_{st})_2V_{24}(U_{s_1t_1})_2(U_{s_2t_2})_4W_{24}(U_{st})_2^\dagger|k\rangle_4]\\
&\quad \otimes[\langle k|_4(U_{st})_2W_{24}^\dagger(U^\dagger_{s_1't_1'})_2(U^\dagger_{s_2't_2'})_4V_{24}^\dagger(T_{st})_2^\dagger|j\rangle_4][U(g)\otimes U(g)]dg|00\rangle\\
&= \frac{1}{n^4(n+1)}\sum_{s_1,t_1}\sum_{s_1',t_1'}\sum_{s_2,t_2}\sum_{s_2',t_2'}\langle\Phi_{s_1t_1}|\chi|\Phi_{s_1't_1'}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s_2't_2'}\rangle\\
&\quad \times\sum_{s,t,j,k,l,l'}\{\mathrm{tr}_2[\langle j|_4(T_{st})_2V_{24}(U_{s_1t_1})_2(U_{s_2t_2})_4|l\rangle_4\langle l|_4W_{24}(U_{st})_2^\dagger|k\rangle_4]\\
&\quad \mathrm{tr}_4[\langle k|_4(U_{st})_2W_{24}^\dagger(U^\dagger_{s_1't_1'})_2(U_{s_2't_2'})_4^\dagger|l'\rangle_4\langle l'|_4V_{24}^\dagger(T_{st})_2^\dagger|j\rangle_4]\\
&\quad +\frac{1}{n^2(n+1)}\sum_{s_1,t_1}\sum_{s_1',t_1'}\sum_{s_2,t_2}\sum_{s_2',t_2'}\langle\Phi_{s_1t_1}|\chi|\Phi_{s_1't_1'}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s_2't_2'}\rangle\\
&\quad \mathrm{tr}_2[(U_{s_1t_1})_2(U^\dagger_{s_1't_1'})_2]\,\mathrm{tr}_4[(U_{s_2t_2})_4(U_{s_2't_2'})^\dagger)_4]\\
&= \frac{1}{n^4(n+1)}\sum_{s_1,t_1}\sum_{s_1',t_1'}\sum_{s_2,t_2}\sum_{s_2',t_2'}\sum_{s,t,j,k}\langle\Phi|_{12}\langle\Phi|_{34}\,\mathrm{tr}_{24}[W_{24}(U_{st})_2^\dagger|k\rangle_4\langle j|_4(T_{st})_2V_{24}(U_{s_1t_1})_2\\
&\quad (U_{s_2t_2})_4](U^\dagger_{s_1t_1})_2(U^\dagger_{s_2t_2})_4\chi_{12}\chi_{34}\,\mathrm{tr}_{24}[V_{24}^\dagger(T_{st})_2^\dagger|j\rangle_4\langle k|_4(U_{st})_2W_{24}^\dagger(U^\dagger_{s_1't_1'})_2(U_{s_2't_2'})^\dagger)_4)]\\
&\quad (U_{s_1't_1'})_2(U_{s_2't_2'})_4|\Phi\rangle_{12}|\Phi\rangle_{34}+\frac{1}{n+1}\\
&= \frac{1}{(n+1)}\sum_{s,t}\langle\Phi|_{12}\langle\Phi|_{34}W_{24}(U_{st})_2^\dagger(T_{st})_2\,\mathrm{tr}_4[V_{24}\chi_{12}\chi_{34}V_{24}^\dagger]\\
&\quad (T_{st})_2^\dagger(U_{st})_2W_{24}^\dagger)|\Phi\rangle_{12}|\Phi\rangle_{34}+\frac{1}{n+1}.
\end{aligned}$$

Then the optimal teleportation fidelity is given by the maximal fidelity of $f_2(\chi)$,

$$\begin{aligned} f_2(\chi)_{max} &= \frac{n^2}{(n+1)} \max_{\Omega,V\in U(n^2)} \{\langle\Phi|_{12}\langle\Phi|_{34}\Omega_{24}\,\mathrm{tr}_4[V_{24}\chi_{12}\chi_{34}V_{24}^\dagger]\Omega_{24}^\dagger|\Phi\rangle_{12}|\Phi\rangle_{34}\} + \frac{1}{n+1} \\ &= \frac{n^2}{(n+1)} \max_{\Omega,V\in U(n^2)} \{\langle\Phi|_{12}\langle\Phi|_{34}\Omega_{13}^T\,\mathrm{tr}_4[V_{24}\chi_{12}\chi_{34}V_{24}^\dagger]\Omega_{13}^*|\Phi\rangle_{12}|\Phi\rangle_{34}\} + \frac{1}{n+1}, \end{aligned}$$

where $\Omega_{24} = W_{24}(U_{st})_2(T_{st})_2$. Rewriting Ω^T as Ω, we get (2). $\Box$

From Theorem 2 we see that the optimal two-channel teleportation fidelity solely depends on the two-channel fully entangled fraction $F_2(\chi)$ of the resource state χ. It can be shown that $F_2(\chi)$ given by (3) is an invariant of local unitary transformations, $\chi_{12}\chi_{34} \to (\mathfrak{U})_{13}(\mathfrak{V})_{24}\chi_{12}\chi_{34}(\mathfrak{U})_{13}^\dagger(\mathfrak{V})_{24}^\dagger$, where $\mathfrak{U}$ and $\mathfrak{V}$ are unitary operators on $H \otimes H$. Theorem 2 also tells us that a resource state χ is useful, namely, it gives better teleportation fidelity than classical channels, if $F_2(\chi) > \frac{1}{n}$.

Let us compare the fidelities between the one-channel and two-channel teleportaion. The usual one-channel teleportaion fidelity $f_1(\chi)_{max}$ is given by [6], $f_1(\chi)_{max} = \frac{nF_1(\chi)}{n+1} + \frac{1}{n+1}$, where $F_1(\chi) = \max_{U\in U(n)}\{\langle\Phi|_{12}U_2^\dagger\chi_{12}U_2|\Phi\rangle_{12}\}$ is the fully entangled fraction. To compare $f_2(\chi)_{max}$ with $f_1(\chi)_{max}$, one only needs to compare $F_2(\chi)$ with $F_1(\chi)$. However, both $F_2(\chi)$ with $F_1(\chi)$ are formidably difficult to calculate analytically. Analytical formulae for $F_1(\chi)$ are only available for some special states χ [21, 22]. Generally one has only estimations of the upper and lower bounds of $F_1(\chi)$ [22, 23]. However, if one takes $W = V$ to be identity, or take $\Omega\, V$ in (3) to be the tensor of two unitary operators $\Upsilon \otimes \Gamma$ with $\Upsilon, \Gamma \in U(n)$, one gets $F_2(\chi) = F_1(\chi)$. Thus the extreme value range of F_2 is larger than that of F_1. Therefore, for any arbitrary state χ, $f_2(\chi)_{max} \geq f_1(\chi)_{max}$, i.e., the two-channel optimal teleportation fidelity is always no less than that of the original one-channel protocol.

In the following, we give numerical calculations of F_2 and F_1 by using the Conjugate Gradient Algorithm [24, 25]. Following the MRPR method introduced in [25], we can get the numerical result of F_1. For F_2, to simplify the computation, we take $V = I_{n\times n}$. Then

$$\begin{aligned} F_2(\chi) &= n \max_{\Omega\in U(n^2)} \{\langle\Phi|_{12}\langle\Phi|_{34}\Omega_{24}\chi_{12}\rho_3\Omega_{24}^\dagger|\Phi\rangle_{12}|\Phi\rangle_{34}\} \\ &= \max_{\Omega\in U(n^2)} \{\sum_{j,j'}\langle\Phi|_{12}\langle j|_4\Omega_{24}\chi_{12}\Omega_{24}^\dagger|j'\rangle_4|\Phi\rangle_{12}\langle j|_3\rho_3|j'\rangle_3\} \\ &= \max_{\Omega\in U(n^2)} \{\sum_{j}\langle\Phi|_{12}\langle j|_3\Omega_{23}\chi_{12}\Omega_{23}^\dagger\rho_3^*|j\rangle_3|\Phi\rangle_{12}\}, \end{aligned}$$

where $\rho_3 = tr_4(\chi_{34})$. Denote $\mathfrak{F}_2(\chi) = \sum_j \langle\Phi|_{12}\langle j|_3\Omega_{23}\chi_{12}\Omega_{23}^\dagger\rho_3^*|j\rangle_3|\Phi\rangle_{12}$. We get $F_2(\chi) = \max_{\Omega_{23}\in U(n^2)} \mathfrak{F}_2$, which is in fact a lower bound of F_2 since we have set $V = I_{n\times n}$. Set $dF = F_2(\chi) - F_1(\chi)$. Fig. 2 shows that for some randomly generated states, one has $F_2(\chi) > F_1(\chi)$.



Figure 2: Hollow triangles for 3-dimension random states, solid triangles for 4-dimension random states.

For the usual one-channel teleportation, it is shown that only resource states χ such that $F_1(\chi) > 1/n$ are useful for teleportation. However, for general states χ, the fully entangled fraction (FEF) $F_1(\chi)$ is quite difficult to compute. Analytical formula of FEF for two-qubit states has been derived by using the method of Lagrange multiplier [21]. The upper bounds of FEF for general high dimensional quantum states have been estimated [23, 26]. Exact results of FEF are also obtained for some special quantum states like isotropic states and Werner states [22]. For a given unknown state, an important issue is to determine whether it is useful for quantum teleportation by experimental measurements. In Ref. [27], the authors show that the set of entangled states which are useful for one-channel quantum teleportation, i.e. their FEFs are great than $1/n$, is convex and compact. The witness operators which detect some entangled states that are useful for teleportation have been derived [28].

From the formulae of the optimal teleportation fidelity $f_2(\chi)_{max}$ (2), it is also obvious that only resource states χ such that $F_2(\chi) > 1/n$ are useful for teleportation. Let $\mathbb{S}$ denote the set of states satisfying $F_2(\chi) \leq \frac{1}{n}$. We have

Theorem 3 $\mathbb{S}$ is convex and compact. There exist witnesses to identify the usefulness of an unknown resource state for two-channel teleportation experimentally.

Proof: (i) The set $\mathbb{S}$ is convex: Let χ_a and $\chi_b \in \mathbb{S}$, namely, $F_2(\chi_a) \leq \frac{1}{n}$, $F_2(\chi_b) \leq \frac{1}{n}$. Consider $\chi_c = \xi\chi_a + (1-\xi)\chi_b$, where $\lambda \in [0,1]$. By the definition of $F_2(\chi) = \max_{\Omega_{23} \in U(n^2)}\{\sum_j \langle\Phi_{00}|_{12}\langle j|_3 \Omega_{23}\chi_{12}\Omega_{23}^{\dagger}\rho_3^*|j\rangle_3|\Phi_{00}\rangle_{12}\}$, We get that $F_2(\chi_c) \leq \xi F_2(\chi_a) + (1-\xi)F_2(\chi_b) \leq \frac{1}{n}$. Thus $\chi_c \in \mathbb{S}$, i.e. $\mathbb{S}$ is convex.

(ii) The set $\mathbb{S}$ is compact: For finite dimensional Hilbert spaces, to show that a set is compact, it is enough to show the set is closed and bounded. $\mathbb{S}$ is bounded since the eigenvalues of χ lies in $[0,1]$ [27]. To see that it is closed, assume that for any two density matrices χ_a and χ_b. The maximal value of $F_2(\chi_a+\chi_b)$ and $F_2(\chi_a)$ are obtained at Ω_{a+b} and Ω_a respectively. Therefore

$$\begin{aligned}
F_2(\chi_a+\chi_b)-F_2(\chi_a) &\leq \sum_j \langle\Phi_{00}|_{12}\langle j|_3(\Omega_{a+b})_{23}(\chi_a)_{12}(\Omega_{a+b})^{\dagger}_{23}tr_4[(\chi_b)_{34}]^*|j\rangle_3|\Phi_{00}\rangle_{12}\\
&+ \sum_j \langle\Phi_{00}|_{12}\langle j|_3(\Omega_{a+b})_{23}(\chi_b)_{12}(\Omega_{a+b})^{\dagger}_{23}tr_4[(\chi_a)_{34}]^*|j\rangle_3|\Phi_{00}\rangle_{12}\\
&+ \sum_j \langle\Phi_{00}|_{12}\langle j|_3(\Omega_{a+b})_{23}(\chi_b)_{12}(\Omega_{a+b})^{\dagger}_{23}tr_4[(\chi_b)_{34}]^*|j\rangle_3|\Phi_{00}\rangle_{12}\\
&\leq \sum_{j,j'} ||\Phi_{00}\rangle||^2||jj'\rangle||^2||\Omega_{a+b}||^2(2||\chi_a||+||\chi_b||)||\chi_b||\\
&= n^2||\Omega_{a+b}||^2(2||\chi_a||+||\chi_b||)||\chi_b||.
\end{aligned}$$

Since the set of all unitary operators is bounded, $||\Omega_{a+b}||^2 \leq v$, where v is a positive real number; $||\chi_a||$ is the maximal eigenvalue of χ_a satisfying $||\chi_a|| \leq 1$. Thus

$$F_2(\chi_a+\chi_b)-F_2(\chi_a) \leq n^2 v(2+||\chi_b||)||\chi_b||.$$

Therefore when $||\chi_b|| \to 0$, $F_2(\chi_a+\chi_b)-F_2(\chi_a) \to 0$, implying that F_2 is a continuous function. Moreover, $F_2(\chi) \in [\frac{1}{n^2},1]$. $F_2=\frac{1}{n^2}$ for $\chi=\frac{1}{n^2}I_{n^2\times n^2}$, and $F_2=1$ for maximally entangled pure states χ. Hence the set $\mathbb{S}=\{\chi:\frac{1}{n^2}\leq F_2(\chi)\leq\frac{1}{n}\}$ is closed.

From the proof above, we can conclude that the set $\mathbb{S}=\{\chi:\frac{1}{n^2}\leq F2(\chi)\leq\frac{1}{n}\}$ is convex and compact. According to the Hahn-Banach theorem [29], any $\chi\notin\mathbb{S}$ can be separated from $\mathbb{S}$ by a hyperplane. This feature enables for the existence of hermitian witness operators and thus experimental ways to detect the usefulness of an unknown state for two-channel teleportation. $\square$

For experimental setup of our two-channel teleportation protocol based on Bell measurement, we give a scheme in terms of photons ($n=2$), see Fig. 3. From spontaneous parametric down-conversion in two adjacent BBO crystals, arbitrary two-photon polarization-entangled pure states can produced [30, 31, 32]. Three entangled pairs (photon 0 and photon p, photon 1 and photon 3, photon 2 and photon 4) can be prepared in state $|\psi\rangle$. Photon 0 is prepared for the initial state while photon p serves as a trigger indicating that a photon to be teleported is under way [33]. The other two pairs play a role as two identical resource pairs. Alice makes a Bell measurement on photon $0,1$. After Bob receives the classical information of measurement results, he performs a corresponding rotation to achieve the maximal teleportation fidelity. The scheme also can be generalized to arbitrary N-channel cases, by simply increasing the number of entangled photon pairs.



Figure 3: Experimental Setup for two-channel photon state teleportation based on Bell measurements. PBS refers to polarizing beam splitter and HWP refers to half wave plate.

Two-channel optimal teleportation protocol based on GHZ measurements

In the above section, we have presented a two-channel teleportation protocol by first making local unitary transformations on the two entangled resources, followed by a standard Bell measurement on the particles 0 and 1. Now we investigate another two-channel teleportation protocol in terms of three-particle Bell-like (GHZ) measurement, see Fig. 4. Alice performs a unitary transformation on particle 1 and 3, then she makes a joint GHZ measurement on her three particles 0, 1 and 3, and delivers the measurement outcomes to Bob by classical means. According to the measurement results, Bob chooses a corresponding unitary transformation T on particle 2 and 4, and trace over the space of particle 4.

First we introduce a set of complete orthogonal normalized generalized GHZ-states $\{|\Phi^s_{rm}\rangle\}$ in $H \otimes H \otimes H$:

$$|\Phi^s_{rm}\rangle = (I \otimes U^s_{rm})|\Phi^0_{00}\rangle,$$

where $|\Phi^0_{00}\rangle = \frac{1}{\sqrt{n}}\sum_{i=0}^{n-1}|iii\rangle$. The unitary matrices $\{U^s_{rm}\}$ are given by: $U^s_{rm} = h^r g^s \otimes h^m$, which satisfy the relations, $tr(U^s_{rm}U^{s'\dagger}_{r'm'}) = n^2\delta_{r,r'}\delta_{m,m'}\delta_{s,s'}$, $U^s_{rm}U^{s\dagger}_{rm} = I_{n^2\times n^2}$. The explicit expressions of the states $\{|\Phi^s_{rm}\rangle\}$ read, $|\Phi^s_{rm}\rangle = \frac{1}{\sqrt{n}}\sum_j w^{js}|j, j+r, j+m\rangle$.

Associated with the states $\{|\Phi^s_{rm}\rangle\}$, one can introduce a set of linear operators $\{\widetilde{U}^{s\dagger}_{rm}\}$: $\widetilde{U}^{s\dagger}_{rm}|i\rangle = \sum_{jk}(U^s_{rm})^*_{ijk}|jk\rangle = \sum_{i',j,k}((U^s_{rm})^*_{i'jk}|j,k\rangle\langle i'|)|i\rangle$, which map $H \to H \otimes H$. It is



Figure 4: Scheme of two-channel optimal teleportation protocol based on GHZ measurements.

easy to show that the correspondence between $\{|\Phi^s_{rm}\rangle\}$ and $\{\widetilde{U}^{s\dagger}_{rm}\}$ is indeed one-one. Explicitly, we have

$$\widetilde{U}^{s\dagger}_{rm} = \sum_j w^{-js}|j+r, j+m\rangle\langle j| = (h^r g^{s*} \otimes h^m)E,$$

where $E = \sum_i |ii\rangle\langle i|$. It is direct to verify that, $\widetilde{U}^s_{rm}\widetilde{U}^{s\dagger}_{rm} = I_{n\times n}$, which maps $H \to H$; and $\widetilde{U}^{s\dagger}_{rm}\widetilde{U}^s_{rm} = (1\otimes h^{m-r})(\sum_j |jj\rangle\langle jj|)(1\otimes h^{m-r})^\dagger$, which maps $H\otimes H \to H\otimes H$.

Theorem 4 The protocol maps an input state ρ_{in} to

$$\begin{aligned}\Lambda^{\{T^s_{rm},W\}}_{\chi^{\otimes 2}}(\rho_{in}) \;=\; & \frac{1}{n^3}\sum_{s_1,t_1}\sum_{s_2,t_2}\sum_{s'_1,t'_1}\sum_{s'_2,t'_2}\langle\Phi_{s_1t_1}|\chi|\Phi_{s'_1t'_1}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s'_2t'_2}\rangle \\ & \times \mathrm{tr}_4\{\sum_{r,m,s}(T^s_{rm})_{24}(U_{s_1t_1})_2(U_{s_2t_2})_4(W)_{24}(\widetilde{U}^{s\dagger}_{rm})_2(\rho_{in})_2 \\ & (\widetilde{U}^s_{rm})_{24}(W^\dagger)_{24}(U_{s'_1t'_1})^\dagger_2(U_{s'_2t'_2})^\dagger_4(T^s_{rm})^\dagger_{24}\}.\end{aligned}$$

Proof. 1). Entangled pure states as resources: with the unknown initial input state given by $|\phi\rangle = \sum_\nu \alpha_\nu|\nu\rangle$, and the two entangled resource state given by $|\Psi\rangle = \sum_{i,j=0}^{n-1}\sum_{k,l=0}^{n-1} a_{ij}|ij\rangle\otimes a_{kl}|kl\rangle$, $\sum_{i,j=0}^{n-1}|a_{ij}|^2 = 1$, applying the operator W on Alice's two resource particles, one has the state $\sum_{i,j=0}^{n-1}\sum_{k,l=0}^{n-1}\sum_{i',k',\nu=0}^{n-1} a_{ij}a_{kl}W^{i'k'}_{ik}\alpha_\nu|\nu i' j k' l\rangle$. From the joint GHZ measurement based on $|\Phi^s_{rm}\rangle$ we get, $\langle\Phi^s_{rm}|(|\phi\rangle\otimes|\Psi'\rangle) = \frac{1}{\sqrt{n}}A_2A_4W_{24}(\widetilde{U}^{s\dagger}_{rm})_2|\phi\rangle_2$. Bob applies a corresponding unitary operator T^s_{rm} on his two resource particles. The resulting state becomes $|\phi\rangle \to \frac{1}{\sqrt{n}}(T^s_{rm})_{24}A_2A_4W_{24}(\widetilde{U}^{s\dagger}_{rm})_2|\phi\rangle_2$. Tracing over the space associated with particle 4, we obtain

$$\begin{aligned}
&\Lambda_{(\chi)}^{\{T^s_{rm},W\}}(\rho_{in}) \\
&= \frac{1}{n}\,\mathrm{tr}_4\{\sum_{r,m,s}(T^s_{rm})_{24}(A)_2(A)_4(W)_{24}(\widetilde{U}^{s\dagger}_{rm})_2(\rho_{in})_2(\widetilde{U}^{s}_{rm})_{24}(W^\dagger)_{24}(A)^\dagger_2(A)^\dagger_4(T^s_{rm})^\dagger_{24}\}.
\end{aligned}$$

2). Arbitrary entangled mixed state as resource: $\sum_{\alpha,\beta} P_\alpha P_\beta |\Psi_{\alpha\beta}\rangle\langle\Psi_{\alpha\beta}|$, where $|\Psi_{\alpha\beta}\rangle = \sum_{i,j=0}^{n-1}\sum_{k,l=0}^{n-1} a^{(\alpha)}_{ij}|ij\rangle \otimes a^{(\beta)}_{kl}|kl\rangle$, $0 \le P_\alpha \le 1$ and $\sum_\alpha P_\alpha = 1$. Following the same procedure as that of the pure resource state case, we have

$$\begin{aligned}
\Lambda_{(\chi)}^{\{T^s_{rm},W\}}(\rho_{in}) &= \frac{1}{n}\sum_{\alpha,\beta} P_\alpha P_\beta\,\mathrm{tr}_4\{\sum_{r,m,s}(T^s_{rm})_{24}(A^{(\alpha)})_2(A^{(\beta)})_4(W)_{24}(\widetilde{U}^{s\dagger}_{rm})_2(\rho_{in})_2 \\
&\quad (\widetilde{U}^{s}_{rm})_{24}(W^\dagger)_{24}(A^{(\alpha)})^\dagger_2(A^{(\beta)})^\dagger_4(T^s_{rm})^\dagger_{24}\}.
\end{aligned}$$

Since $A^{(\alpha)} = \sum_{s,t} a^{(\alpha)}_{st} U_{st}$ and $n\sum_\alpha p_\alpha a^{(\alpha)}_{st} a^{(\alpha)*}_{s't'} = \langle\Phi_{st}|\chi|\Phi_{s't'}\rangle$, we get

$$\begin{aligned}
\Lambda_{(\chi)}^{\{T^s_{rm},W\}}(\rho_{in}) &= \frac{1}{n^3}\sum_{s_1,t_1}\sum_{s_2,t_2}\sum_{s'_1,t'_1}\sum_{s'_2,t'_2}\langle\Phi_{s_1t_1}|\chi|\Phi_{s'_1t'_1}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s'_2t'_2}\rangle \\
&\times\ \mathrm{tr}_4\{\sum_{r,m,s}(T^s_{rm})_{24}(U_{s_1t_1})_2(U_{s_2t_2})_4(W)_{24}(\widetilde{U}^{s\dagger}_{rm})_2(\rho_{in})_2 \\
&\quad (\widetilde{U}^{s}_{rm})_{24}(W^\dagger)_{24}(U_{s'_1t'_1})^\dagger_2(U_{s'_2t'_2})^\dagger_4(T^s_{rm})^\dagger_{24}\},
\end{aligned}$$

which is valid also for any mixed input states ρ_{in}. □

It is direct to prove that the quantum channel $\Lambda_{\chi^{\otimes 2}}^{\{T^s_{rm},W\}}$ is trace preserving, namely, $\mathrm{tr}[\Lambda_{(\chi)}^{(\{T^s_{rm},W\})}(\rho_{in})] = 1$

Theorem 5 The optimal teleportaion fidelity of the two-channel protocol based on GHZ measurements is given by

$$f'_{2\max}(\chi) = \frac{nF'_2(\chi)}{(n+1)} + \frac{1}{n+1}, \tag{4}$$

where

$$\begin{aligned}
F'_2(\chi) &= \frac{1}{n}\max_{\{W,T^s_{rm}\}}\sum_{r,m,s,i}\{\langle\Phi|_{12}\langle\Phi|_{34}[W_{24}(h^r g^{s*})_2(h^m)_4(E_i)_{24}(T^s_{rm})_{24}](\chi_{12}\chi_{34}) \\
&\quad [(T^{s\dagger}_{rm})_{24}(E^\dagger_i)_{24}(h^r g^{s*})^\dagger_2(h^m)^\dagger_4 W^\dagger_{24}]|\Phi\rangle_{12}|\Phi\rangle_{34}\}
\end{aligned}$$

and $E_i = \sum_{j=0}^{n-1}|jj\rangle\langle ji|$.

Proof. Using the Schur's lemma and averaging over all possible input states, we have

$$\begin{aligned}
f'(\chi) &= \overline{\langle\phi_{in}|\Lambda^{\{T^s_{rm},W\}}_{(\chi)}(\rho_{in})|\phi_{in}\rangle} = \frac{1}{n^3}\sum_{s_1,t_1}\sum_{s_2,t_2}\sum_{s'_1,t'_1}\sum_{s'_2,t'_2}\langle\Phi_{s_1t_1}|\chi|\Phi_{s'_1t'_1}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s'_2t'_2}\rangle \\
&\times \sum_{r,m,s,i}\langle 00|\int_G[U(g)^\dagger\otimes U(g)^\dagger]\langle i|_4[(T^s_{rm})_{24}(U_{s_1t_1})_2(U_{s_2t_2})_4(W)_{24}(\widetilde{U}^{s\dagger}_{rm})_2] \\
&\otimes [(\widetilde{U}^s_{rm})_{24}(W^\dagger)_{24}(U^\dagger_{s'_1t'_1})_2(U^\dagger_{s'_2t'_2})_4(T^{s\dagger}_{rm})_{24}]|i\rangle_4[U(g)\otimes U(g)]dg|00\rangle \\
&= \frac{1}{n^4(n+1)}\sum_{s_1,t_1}\sum_{s_2,t_2}\sum_{s'_1,t'_1}\sum_{s'_2,t'_2}\langle\Phi_{s_1t_1}|\chi|\Phi_{s'_1t'_1}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s'_2t'_2}\rangle \\
&\times \sum_{r,m,s,i}\{\mathrm{tr}_{24}[(T^s_{rm})_{24}(U_{s_1t_1})_2(U_{s_2t_2})_4(W)_{24}(\widetilde{U}^{s\dagger}_{rm})_2\langle i|_4] \\
&\quad \mathrm{tr}_{24}[|i\rangle_4(\widetilde{U}^s_{rm})_{24}W^\dagger_{24}(U^\dagger_{s'_1t'_1})_2(U^\dagger_{s'_2t'_2})_4(T^{s\dagger}_{rm})_{24}] \\
&+ \mathrm{tr}_2[\langle i|_4(T^s_{rm})_{24}(U_{s_1t_1})_2(U_{s_2t_2})_4(W)_{24}(\widetilde{U}^{s\dagger}_{rm})_2 \\
&\quad (\widetilde{U}^s_{rm})_{24}(W^\dagger)_{24}(U^\dagger_{s'_1t'_1})_2(U^\dagger_{s'_2t'_2})_4(T^{s\dagger}_{rm})_{24}|i\rangle_2]\}.
\end{aligned}$$

From the following identity, $\sum_{r,m,s}\widetilde{U}^{s\dagger}_{rm}\,A\,\widetilde{U}^s_{rm} = n\,tr(A)\,I_d\otimes I_d$, one gets

$$\begin{aligned}
f'(\chi) &= \frac{1}{n^4(n+1)}\sum_{s_1,t_1}\sum_{s_2,t_2}\sum_{s'_1,t'_1}\sum_{s'_2,t'_2}\sum_{r,m,s,i}\{\langle\Phi|_{12}\langle\Phi|_{34}[(U^\dagger_{s_1t_1})_2(U^\dagger_{s_2t_2})_4] \\
&\quad \mathrm{tr}_{24}[(U_{s_1t_1})_2(U_{s_2t_2})_4W_{24}(h^rg^{s*})_2(h^m)_4(E_i)_{24}(T^s_{rm})_{24}](\chi_{12}\chi_{34}) \\
&\quad \mathrm{tr}_{24}[(T^{s\dagger}_{rm})_{24}(E^\dagger_i)_{24}(h^rg^{s*})^\dagger_2(h^m)^\dagger_4W^\dagger_{24}(U^\dagger_{s'_1t'_1})_2(U^\dagger_{s'_2t'_2})_4][(U_{s'_1t'_1})_2(U_{s'_2t'_2})_4]|\Phi\rangle_{12}|\Phi\rangle_{34} \\
&+ \frac{1}{n^2(n+1)}\sum_{s_1,t_1}\sum_{s_2,t_2}\sum_{s'_1,t'_1}\sum_{s'_2,t'_2}\langle\Phi_{s_1t_1}|\chi|\Phi_{s'_1t'_1}\rangle\langle\Phi_{s_2t_2}|\chi|\Phi_{s'_2t'_2}\rangle\ \mathrm{tr}\{U_{s_1t_1}U^\dagger_{s'_1t'_1}\}\ \mathrm{tr}\{U_{s_2t_2}U^\dagger_{s'_2t'_2}\} \\
&= \frac{1}{(n+1)}\sum_{r,m,s,i}\{\langle\Phi|_{12}\langle\Phi|_{34}[W_{24}(h^rg^{s*})_2(h^m)_4(E_i)_{24}(T^s_{rm})_{24}](\chi_{12}\chi_{34}) \\
&\quad [(T^{s\dagger}_{rm})_{24}(E^\dagger_i)_{24}(h^rg^{s*})^\dagger_2(h^m)^\dagger_4W^\dagger_{24}]|\Phi\rangle_{12}|\Phi\rangle_{34}\} + \frac{1}{n+1}.
\end{aligned}$$

Thus the optimal teleportation fidelity $f'_{2\,\max}(\chi)$ is then given by (4).

It can be shown that the TFEF $F'_2(\chi)$ is also an invariant under local unitary transformations on the resource pairs. The Experimental scheme of the two-channel optimal teleportation protocol based on GHZ measurements[34] is shown in Fig. 5.

Conclusions and Discussions

We have proposed two general two-channel quantum teleportation protocols, the corresponding optimal teleportation fidelities have been explicitly derived. It turns out that



Figure 5: Experimental Setup for two-channel photon state teleportation based on GHZ measurements. PBS refers to polarizing beam splitter, HWP refers to half wave plate, and SPR refers to single-qubit polarization rotations.

the optimal teleportation fidelities only depend on the two-channel fully entangled fractions, which are invariants under local unitary transformations on the resource states. It has been shown that the two-channel protocol based on Bell measurement can increase the teleportation fidelity bypassing the usual one-channel protocol. To get better and better teleportation fidelity, avoid complicated distillation procedure and save entangled resources at the same time, one may generalize the two-channel teleportation protocol to multi-channel ones.

Acknowledgement

Financial support from the National Natural Science Foundation of China (Grant Nos. 11275131, 11375141, 11434013, 11425522) and the ”973” program (2010CB922904) are gratefully acknowledged. Drs. Huangjun Zhu and Xiaolong Du helped us a lot on the numerical algorithm. We also would like to thank Profs. Kai Chen, Zhihua Chen, Ming Li, Drs. Teng Ma and Bin Chen for their helpful discussions.

References

[1] Nielsen, M. A. & Chuang, I. L. *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England, 2000).

[2] Bennett, C. H. et al. Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels. *Phys. Rev. Lett.* **70**, 1895 (1993).

[3] D'Ariano, G. M., Lo Presti, P. & Sacchi, M. F. Bell measurements and observables. *Phys. Lett. A* **272**, 32 (2000).

[4] Bowen, G. & Bose, S. Teleportation as a Depolarizing Quantum Channel, Relative Entropy, and Classical Capacity. *Phys. Rev. Lett.* **87**, 267901 (2001).

[5] Albeverio, S., Fei, S. M. & Yang, W. L., Teleportation with an Arbitrary Mixed Resource as a Trace-Preserving Quantum Channel. *Commun. Theor. Phys.* **38**, 301-304 (2002).

[6] Albeverio, S., Fei, S. M. & Yang, W. L., Optimal teleportation based on bell measurements. *Phys. Rev. A* **66**, 012301 (2002).

[7] Alber, G. et al. *Quantum Information: An Introduction to Basic Concepts and Experiments* (Springer Tracts in Modern Physics, 2001).

[8] Abeyesinghe, A., Devetak, I., Hayden, P. & Winter, A. The mother of all protocols: restructuring quantum informations family tree. *Proc. R. Soc. A* **465(2108)**: 2537-2563 (2009).

[9] Bennett, C. H. et al., Purification of Noisy Entanglement and Faithful Teleportation via Noisy Channels. *Phys. Rev. Lett.* **76**, 722-725 (1996).

[10] Bennett, C. H., Bernstein, H. J., Popescu, S. & Schumacher, B. Concentrating partial entanglement by local operations. *Phys. Rev.A* **53**, 2046 (1996).

[11] Horodecki, M., Horodecki, P. & Horodecki, R. Inseparable Two Spin- 12 Density Matrices Can Be Distilled to a Singlet Form. *Phys. Rev. Lett.* **78** 574 (1997).

[12] Horodecki, M., Horodecki, P. & Horodecki, R. Mixed-State Entanglement and Distillation: Is there a Bound Entanglement in Nature? *Phys. Rev. Lett.* **80** 5239 (1998).

[13] Peres, A. Separability Criterion for Density Matrices. *Phys. Rev. Lett.* **77** 1413 (1996).

[14] Horodecki, R., Horodecki, P., Horodecki, M. & Horodecki, K. Quantum entanglement. *Rev, Mod. Phys.* **81**, 865 (2009).

[15] Horodecki, P. Separability criterion and inseparable mixed states with positive partial transposition. *Phys.Lett. A* **232** 333(1996).

[16] Horodecki, P., Horodecki, M. & Horodecki, R. Bound Entanglement Can Be Activated. *Phys.Rev.Lett.* **82** 1056-1059 (1999).

[17] Masanes, L. All Bipartite Entangled States Are Useful for Information Processing. *Phys. Rev. Lett.* **96** 150501 (2006).

[18] Cavalcanti, D., Acin, A., Brunner, N. & Vertesi, T. All quantum states useful for teleportation are nonlocal resources. *Phys. Rev. A* **87** 042104 (2013).

[19] Vrtesi, T. & Brunner, N. Disproving the Peres conjecture: Bell nonlocality from bipartite bound entanglement. *Nature Communications* **5** 5297 (2014).

[20] Werner, R. F. All teleportation and dense coding schemes. *J. Phys. A: Math. Gen.* **34** 7081(2001).

[21] Grondalski, J., Etlinger, D. M. & James, D. F. V. The fully entangled fraction as an inclusive measure of entanglement applications. *Phys. Lett. A* **300**, 573 (2002).

[22] Zhao, M. J., Li, Z. G., Fei, S. M. & Wang, Z. X. A Note on Fully Entangled Fraction. *J. Phys. A* **43**, Math. Theor. 275203 (2010).

[23] Li, M., Fei, S. M. & Wang, Z. X. Upper bound of the fully entangled fraction. *Phys. Rev. A* **78**, 032332 (2008).

[24] Shrwchuk, J. R. *An Introduction to the Conjugate Gridient Method Without the Agonizing Pain*, (August 1994).

[25] Zhang, Li., Zhou, W. J. & Li, D. H. A descent modified PolakCRibi‘ereCPolyak conjugate gradient method and its global convergence. *Journal of Numerical analysis* **26** 629-640 (2006).

[26] Gu, R. J., Li, M., Fei, S. M. & Li-Jost, X. On Estimation of Fully Entangled Fraction. *Commun. Theor. Phys.* **53**, 265 (2010).

[27] Ganguly, N., Adhikari, S., Majumdar, A. S. & Chatterjee, J. Entanglement Witness Operator for Quantum Teleportation. *Phys. Rev. Lett.* **107**, 270501 (2011).

[28] Zhao, M. J., Fei, S. M. & Li-Jost, X. Complete entanglement witness for quantum teleportation. *Phys. Rev. A* **85**, 054301 (2012).

[29] Holmes, R. B. *Geometric Functional Analysis and its Applications*, (Springer-Verlag 1975).

[30] Kwiat, P. G., Mattle, K., Weinfurter, H. & Zeilinger, A. New High-Intensity Source of Polarization-Entangled Photon Pairs. *Phys. Rev. Lett.* **75**, 4337 (1995).

[31] White, A. G., James, D. F. V., Eberhard P. H., & Kwiat P. G. Nonmaximally Entangled States: Production, Characterization, and Utilization. *Phys. Rev. Lett.* **83**, 3103 (1999).

[32] Zhang, C. Preparation of polarization-entangled mixed states of two photons. *Phy. Rev. A* **69**, 014304(2004).

[33] Bouwmeester, D. et al. Experimental quantum teleportation *Nature* (London) **390**, 575-579 (1997).

[34] Pan J. W. & Zeilinger, A. Greenberger-Horne-Zeilinger-state analyzer. *Phys. Rev. A* **57**, 2208-2211 (1998).